

Parsons (*loc. cit.*) states that this species has been collected on *Yucca elata*. Saguaro (*Cereus giganteus*) is certainly the principal host plant in the Tucson area. We have never taken the species in other blossoms. *C. pallipennis* frequents saguaro blossoms only in small numbers.

---

*EMELINUS MELSHEIMERI* (LEC.) IN ARIZONA (COLEOPTERA: ADERIDAE).—This species has not been recorded from west of Illinois and would be expected to occur only in the mesophytic portions of the Northeast and Midwest. Leng's record from Florida may refer to a misdetermined *E. ashmeadi*. A single male, collected at light by Anthony Ross, of the Department of Entomology, University of Arizona, in Pinery Canyon, W. slope Chiricahua Mts., July 11, 1958, agrees perfectly with the male from Illinois mentioned previously (Werner, 1956, *Psyche* 63: 32). This brings the total of described species known from Arizona to three. A small portion of the fauna of the Chiricahua Mts. appears to have been derived from that of the Midwest or at least from the same source. J. W. Green has recently identified a specimen of the lycid *Caenia dimidiata* (Fab.) from Pinery Canyon, July 9, 1955, Butler and Werner collectors. He expressed some surprise at its occurrence there, since it was previously known from only as far west as Arkansas. — F. G. WERNER, University of Arizona.